arXiv:1109.3894v3 [quant-ph] 20 Apr 2012

Exact Solution of Schrödinger Equation with Three Physical Potentials Using the Nikiforov-Uvarov Method

Falaye, B. J.[1] , **Oyewumi, K. J.**[2] and **Abbas, M.**[3]

Theoretical Physics Section, Department of Physics
University of Ilorin, P. M. B. 1515, Ilorin, Nigeria.

Abstract

In this paper we present the exact solution of one dimensional Schrödinger equation for Woods-Saxon plus Rosen-Morse plus symmetrical double well potential via Nikiforov-Uvarov mathematical method. The eigenvalues and eigenfunctions of this potential are obtained. The energy equations and the corresponding wave function for special cases of this potential are briefly discussed. The PT-symmetry and Hermiticity for this potential are also considered.

Keywords: Schrödinger equation, Woods-Saxon, Rosen-Morse, symmetrical double well potential, Nikiforov-Uvarov

PACs No. 03.65.Ge; 03.65.Pm

1 Introduction

It is well known that the exact solutions of the Schrödinger equation for some physical potential, plays an essential role in non-relativity quantum mechanics and are possible only for a few set of quantum systems.

In recent time, a number of researchers have studied the solution of the Schrödinger equation. For example, these investigation have employed Woods-Saxon Potential [1, 2, 3, 4], Harmonic oscillator potential [5], Hulthén potential [6, 7, 8], Rosen-Morse potential [9, 10], Manning-Rosen potential [11] etc. Several methods such as the Nikivorov-Uvarov method, Supper-Symmetric quantum mechanics, Factorization method, etc, have been used to solve differential equations arising from these considerations.

In this present work, our focus is to solve the one dimensional Schrödinger equation for Woods-Saxon plus Rosen-Morse plus symmetrical double well potential using the

---

[1] E-mail: fbjames11@physicist.net
[2] E-Mail: kjoyewumi66@unilorin.edu.ng
[3] E-Mail: abbasmustapha1@yahoo.com

Nikiforov-Uvarov mathematical method. The results include the PT-symmetric and the non PT-symmetric solution of this potential

The paper is organized as follows, section 2; we briefly review Nikiforov-Uvarov $(N-U)$ Mathematical method. In section 3, by using this method, we obtain the energy eigenvalue and eigenfunction for real and complex cases of this potential. We investigate the PT-symmetric and non PT-symmetric of this potential in sect. 4 and sect. 5. In section 6, we investigate the special cases of this potential and section 7 devoted to conclusion.

2 The Nikiforov-Uvarov (N-U) Method

The N-U method is based on solving a second order linear differential equation by reducing it to a generalized equation of hypergeometric type [12, 13]. By introducing an appropriate coordinate transformation $z = z(r)$, this equation can be re-written in the following form.

$$\Psi''(z) + \frac{\tilde{\tau}(z)}{\sigma(z)}\Psi'(z) + \frac{\tilde{\sigma}(z)}{\sigma^2(z)}\Psi(z) = 0 \tag{1}$$

where $\sigma(z)$ and $\tilde{\sigma}(z)$ are polynomials, at most of second degree, and $\tilde{\tau}(z)$ is a first degree polynomial. Now, if one takes the following factorization

$$\Psi(z) = y(z)\phi(z). \tag{2}$$

The equation (1) reduces to a hypergeometric type equation

$$\sigma(z)y''(z) + \tau(z)y'(z) + \lambda y(z) = 0, \tag{3}$$

where

$$\tau(z) = \tilde{\tau}(z) + 2\pi(z). \tag{4}$$

The function π and the parameter λ require for this method are define as follows

$$\pi(z) = \left(\frac{\sigma'(z) - \tilde{\tau}(z)}{2}\right) \pm \sqrt{\left(\frac{\sigma'(z) - \tilde{\tau}(z)}{2}\right)^2 + \tilde{\sigma}(z) + k\sigma(z)} \tag{5}$$

and

$$\lambda = k + \pi'(z) \tag{6}$$

In other to find the value of k, the expression under the square root must be a square of a polynomial. This gives the polynomial $\pi(z)$ which is dependent on the transformation function $z(r)$. Also the parameter λ defined in equation (6) takes the form

$$\lambda = \lambda_n = -n\tau' - \left[\frac{n(n-1)}{2}\sigma''\right]. \tag{7}$$

The polynomial solutions $y_n(z)$ are given by the Rodrigue relation

$$y_n(z) = \frac{N_n}{\rho(z)}\frac{d^n}{dz^n}\left[\sigma^n(z)\rho(z)\right] \tag{8}$$

where N_n is the normalization constant and $\rho(z)$ is the weight function satisfying

$$\frac{d}{dz}\left[\sigma(z)\rho(z)\right] = \tau(z)\rho(z). \tag{9}$$

Second part of the wave function $\phi(z)$ can be obtained from the relation

$$\pi(z) = \sigma(z)\frac{d}{dz}[\ln\phi(z)] \tag{10}$$

3 Bound state solutions

The one dimensional time independent Schrödinger Equation can be written as

$$\frac{d^2\Psi(x)}{dx^2} + \left[E - V(x)\right]\Psi(x) = 0 \quad , \quad for \;\; \hbar = 2\mu = 1 \tag{11}$$

The generalized Woods-Saxon plus Rosen-Morse plus generalized symmetrical double well Potential can be expressed as

$$V(x) = \left\{ \begin{array}{c} \frac{V_1 e^{-2\alpha x}}{1+qe^{-2\alpha x}} + \frac{V_2 e^{-4\alpha x}}{(1+qe^{-2\alpha x})^2} + V_3 \sec h_q^2(\alpha x) \\ +V_4 \tan h_q(\alpha x) - V_5 \tan h_q^2(\alpha x) + V_6 \sec h_q^2(\alpha x) \end{array} \right\} \tag{12}$$

Using the following deformed hyperbolic function [14] we have

$$\sin h_q x = \frac{e^x - qe^{-x}}{2}, \cos h_q x = \frac{e^x + qe^{-x}}{2}, \tan h_q x = \frac{\sin h_q x}{\cos h_q x}$$

$$\sec h_q x = \frac{1}{\cos h_q x}, cosech_q x = \frac{1}{\sin h_q x}, \cot h_q x = \frac{\cos h_q x}{\sin h_q x} \tag{13}$$

By using

$$z = -e^{-2\alpha x} \tag{14}$$

We obtained

$$\frac{d^2}{dx^2} = 4\alpha^2 z^2 \frac{d^2}{dz^2} + 4\alpha z\frac{d}{dz} \tag{15}$$

Putting Eqs. (15), (14), (12) and appropriate parameter in Eq. (13) into Eq. (11), we have

$$\Psi^{''}(z) + \frac{1-qz}{z(1-qz)}\Psi^{'}(z) + \frac{1}{z^2(1-qz)^2}[z^2\left\{\beta_1 q - \beta_2 - q^2(\beta_4+\beta_5+\xi)\right\} + z\left\{-\beta_1 - 4(\beta_3+\beta_6) - 2q\beta_5 + 2q\xi\right\} + (\beta_4 - \xi)] \tag{16}$$

Where we have used the following notations

$$\xi = \frac{-E}{4\alpha^2} \quad and \quad \beta_i = \frac{V_i}{4\alpha^2} \;\;, \;\; i = 1-6 \tag{17}$$

Comparing Eq. (16) with Eq. (1), we have

$$\sigma(z) = z^2\left\{\beta_1 q - \beta_2 - q^2(\beta_4+\beta_5+\xi)\right\} + z\left\{-\beta_1 - 4(\beta_3+\beta_6) - 2q\beta_5 + 2q\xi\right\} + (\beta_4 - \xi)$$

$$\tilde{\tau}(z) = 1 - qz \;, \;\; \sigma(z) = z - qz^2 \tag{18}$$

Inserting the values of these parameters with $\sigma'(z) = 1 - 2qz$ into Eq. (5), we obtained

$$\pi(z) = -\frac{qz}{2} \pm \frac{1}{2}\left[\{2\delta - \eta H\}qz - 2\delta\right], \quad for \;\; k = 2q(\beta_4 - \beta_5) - 4(\beta_3 + \beta_6) - \beta_1 + \eta H q\delta \tag{19}$$

Where we have used the following notations

$$\eta = \pm 1, \quad \delta = \sqrt{\xi - \beta_4}, H = \sqrt{\frac{4\beta_2}{q^2} + 16\frac{(\beta_3+\beta_6)}{q} + 12\beta_5 + 1} \tag{20}$$

By choosing an appropriate parameter of k in $\pi(z)$ which satisfies the condition $\tilde{\tau}(z) < 0$, one gets

$$\pi(z) = -\frac{qz}{2} - \frac{1}{2}\left[\{\delta - \eta H\}qz - 2\delta\right] \quad for \;\; k = 2q(\beta_4 - \beta_5) - 4(\beta_3 + \beta_6) - \beta_1 + \eta H q\delta \tag{21}$$

From Eq. (6), we obtained the parameter λ as

$$\lambda = 2q(\beta_4 - \beta_5) - 4(\beta_3 + \beta_6) - \beta_1 + \eta H q\delta - \frac{q}{2} - \frac{1}{2}\{\{2\delta - \eta H\}qz\} \tag{22}$$

Also we obtain the parameter λ from Eq. (7) as

$$\lambda = \lambda_n = nq\{(n+1) + 2\delta - \eta H)\} \tag{23}$$

Comparing Eqs. (22) and (23), we obtained the exact energy eigenvalues equation as

$$E_n = -\frac{\alpha^2}{4}\left[\frac{\frac{2qV_4+qV_5-V_1}{q\alpha^2} + \frac{V_2}{q^2\alpha^2}}{2n+1-\eta\sqrt{\frac{V_2}{q^2\alpha^2} + 4\frac{(V_3+V_6)}{q\alpha^2} + 3\frac{V_5}{\alpha^2} + 1}} - \left(2n+1-\eta\sqrt{\frac{V_2}{q^2\alpha^2} + 4\frac{(V_3+V_6)}{q\alpha^2} + 3\frac{V_5}{\alpha^2} + 1}\right)\right]^2 - V_4$$

$$n \geq 0, q \geq 1 \tag{24}$$

Where we have substituted for $\beta_1,\ \beta_2,\ \beta_3,\ \beta_4,\ \beta_5,\ \beta_6,\ \delta$ and H. From Eq. (9) it can be shown that the weight function $\rho(z)$ is

$$\rho(z) = z^{2\delta}(1-qz)^{-\eta H} \tag{25}$$

Substituting Eq. (25) into the Rodrigue relation (8), we obtained

$$y_n(z) = N_n P_n^{(2\delta,-\eta H)}(1-2qz) \tag{26}$$

Where, we have used the properties of Jacobi Polynomial [15, 16]. We obtain the other part of the wave function from Eq. (9) as

$$\phi\left(z\right) = z^{\delta}(1-qz)^{\frac{1}{2}(1-\eta H)} \tag{27}$$

Finally we obtained the wave function $\Psi_n\left(z\right) = \phi\left(z\right) y_n\left(z\right)$ as

$$\Psi_n(z) = N_n z^{\sqrt{\xi-\beta_4}}(1-qz)^{\frac{1}{2}\left(1-\eta\sqrt{\frac{V_2}{q^2\alpha^2}+4\frac{(V_3+V_6)}{q\alpha^2}+3\frac{V_5}{\alpha^2}+1}\right)} P_n^{\left(2\sqrt{\xi-\beta_4},-\eta\sqrt{\frac{V_2}{q^2\alpha^2}+4\frac{(V_3+V_6)}{q\alpha^2}+3\frac{V_5}{\alpha^2}+1}\right)}(1-2qz) \tag{28}$$

Where N_n is the normalization factor to be determined from normalization condition

$$\int_0^1 \left|\Psi_n\left(z\right)\right|^2 dz = 1 \tag{29}$$

This can further be written as

$$N_n^2 \int_0^1 z^{2c}(1-qz)^{(1+2d)}\left[P_n^{(2c,2d)}(z)\right]^2 = 1 \tag{30}$$

Where $c = \delta$ and $d = -\frac{\eta H}{2}$. Using the different forms of Jacobi polynomial [15, 16]

$$P_n^{(c,d)}\left(z\right) = 2^{-n}\sum_{r=0}^{n}(-1)^n \binom{n+c}{p}\binom{n+d}{n-p}(1-z)^{n-p}(1-z)^p \tag{31}$$

$$P_n^{(c,d)}\left(z\right) = \frac{(\Gamma n+c+1)}{n!\Gamma(n+c+d+1)}\sum_{r=0}^{n}\binom{n}{r}\frac{\Gamma(n+c+d+r+1)}{r+c+1}\left(\frac{z-1}{2}\right)^r \tag{32}$$

Where $\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{\Gamma(n+1)}{\Gamma(r+1)\Gamma(n-r+1)}$

We obtained $\left[P_n^{(2c,2d)}(x)\right]^2$ from Eqs. (31) and (32) as

$$\left[P_n^{(2c,2d)}(z)\right]^2 = z^{n+r-p}(1-qz)^p A_{nq}\left(p,r\right) \tag{33}$$

Putting Eq. (33) into (30), we have

$$N_n^2 A_n q\left(p,r\right)\int_0^1 z^{n+r-p+2c}(1-qz)^{1+2d+p}dz=1 \tag{34}$$

Before presenting the normalization constant N_n, let us recall a relation of hypergeometric function, which is used to solve Eq. (34) and present N_n

$$\int_1^0 z^{c-a-1}(1-qz)^{-b}ds={}_2F_1(a,b;a+1;q)\times\frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \tag{35}$$

$$\text{Provided } Re(c)<Re(a)>0,|\arg(1-q)|<\pi$$

$$\int_0^1 z^{a-1}(1-qz)^{-b}ds=\frac{1}{a}[{}_2F_1(a,b;a+1;q)] \tag{36}$$

Using the above Eq. (36), we obtain the normalization factor N_n as

$$N_n=\sqrt{\frac{B_{nq}(p,r)}{A_{nq}\left(p,r\right)}} \tag{37}$$

Where

$$A_{nq}(p,r)=\frac{(-1)^n[\Gamma\left(n+2c+1\right)]^2\Gamma(n+2d+1)}{\Gamma\left(n+2c-p+1\right)\Gamma\left(r+2d+1\right)\Gamma(2c+2d+1)}\sum_{n}^{P=0}\frac{(-1)^{p+r}q^{n-p+r}\Gamma(n+2c+2d+r+1)}{p!r!\left(n-p\right)!\left(n-r\right)!\Gamma(p+2d+1)} \tag{38}$$

$$(B_{nq}(p,r))^{-1}=\frac{1}{n+r-p+2\delta+1}{}_2F_1(n+r-p+2\delta+1,\eta H-1-p;n+r-p+2\delta+1;q) \tag{39}$$

$$c=\delta,\quad \delta=\sqrt{\xi-\beta_4},\quad \beta_4=\frac{V_4}{4\alpha^2},\quad d=-\frac{\eta H}{2},\quad H=\sqrt{\frac{V_2}{q^2\alpha^2}+4\frac{(V_3+V_6)}{q\alpha^2}+3\frac{V_5}{\alpha^2}+1} \tag{40}$$

4 PT -Symmetry and non-Hermiticity

When a Hamiltonian commutes with PT, it is called PT-symmetric Hamiltonian, i.e. $[H,PT]=0$, where the parity operator P and the time reversal operator T, satisfying the following relations.

$$PpP^{-1}=-P=TpT^{-1},\quad PxP^{-1}=-x,\quad TAT^{-1}=-iA \tag{41}$$

In this case, we set the potentials parameter in Eq. (12) as $V_1,\ V_2,\ V_3,\ V_4,\ V_5,\ V_6,$ $q\in\Im$ and $\alpha\in\cap\Im\,(\alpha\to i\alpha)$. Where $\Im$ and $\cap\Im$ denotes the set of purely real numbers and purely complex numbers respectively.

This replacement and after some manipulations in Eq. (12) leads to the following potential

$$V(x)=\frac{\left\{\begin{array}{c}-V_1\left[1+q(\cos 2\alpha x-isin2\alpha x)\right]+V_2\left[\cos 2\alpha x-isin2\alpha x\right]-4(V_3+V_6)\\ -V_4\left[(1-q^2)\cos 2\alpha x+i(1+q^2)\sin 2\alpha x\right]+V_5\left[(1+q^2)\cos 2\alpha x+i(1-q^2)\sin 2\alpha x-2q\right]\end{array}\right\}}{(1+q^2)\cos 2\alpha x+(1-q^2)\,isin2\alpha x+2q} \tag{42}$$

Using Eqs. (11) and (42) and making the corresponding parameter replacements in Eqs. (24) and (28), we obtained energy eigenvalues equation and eigenfunction respectively as

$$E_n=\frac{\alpha^2}{4}\left[\frac{\frac{V_1-2qV_4-qV_5}{q\alpha^2}-\frac{V_2}{q^2\alpha^2}}{2n+1-\eta\sqrt{1-\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}}-\left(2n+1-\eta\sqrt{1-\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)\right]^2-V_4$$

$$n\geq 0, q\geq 1 \tag{43}$$

$$\Psi_n\left(z\right)=C_n z^{\sqrt{\xi-\beta_4}}(1-qz)^{\frac{1}{2}\left(1-\eta\sqrt{1-\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)}P_n^{\left(2\sqrt{\xi-\beta_4},-\eta\sqrt{1-\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)}(1-2qz) \tag{44}$$

Using the above Eq.(44), we can consequently obtain the normalization constant C_n as

$$C_n=\sqrt{\frac{B_{nq}(p,r)}{A_{nq}(p,r)}} \tag{45}$$

Where

$$A_{nq}(p,r)=\frac{(-1)^n[\Gamma\left(n+2c+1\right)]^2\Gamma(n+2d+1)}{\Gamma\left(n+2c-p+1\right)\Gamma\left(r+2d+1\right)\Gamma(2c+2d+1)}\sum_{n}^{P=0}\frac{(-1)^{p+r}q^{n-p+r}\Gamma(n+2c+2d+r+1)}{p!r!\left(n-p\right)!\left(n-r\right)!\Gamma(p+2d+1)} \tag{46}$$

$$\left(B_{nq}\left(p,r\right)\right)^{-1}=\frac{1}{n+r-p+2\delta+1}{}_2F_1\left(n+r-p+2\delta+1,\eta H-1-p;n+r-p+2\delta+1;q\right) \tag{47}$$

$$c=\delta=\sqrt{\xi-\beta_4},\beta_4=\frac{V_4}{4\alpha^2},d=-\frac{\eta H}{2},H=\sqrt{1-\frac{V_2}{q^2\alpha^2}-4\frac{V_3+V_6}{q\alpha^2}-3\frac{V_5}{\alpha^2}}$$

5 Non PT -symmetry and non Hermiticity

We obtain another form of the potential by assuming some of the parameters as pure imaginary,i.e. $q\to iq$, $\alpha\to i\alpha$, $V_1\to iV_1$, $V_3\to iV_3$, $V_6\to iV_6$. Under this replacement and after some manipulations, the potential takes the form

$$V\left(x\right)=\frac{\left\{\begin{array}{c}V_1\left[qcos\alpha x+i\left(1+qcos2\alpha x\right)\right]+V_2[\cos 2\alpha x-isin2\alpha x]-i(V_3+V_6)\\ -V_4\left[(1+q^2)\cos 2\alpha x+i(1-q^2)\sin 2\alpha x\right]+V_5\left[\left(1+q^2\right)\cos 2\alpha x+i\left(1-q^2\right)\sin 2\alpha x-2iq\right]\end{array}\right\}}{\left(1-q^2\right)\cos 2\alpha x+i\left(1+q^2\right)\sin 2\alpha x+2iq} \tag{48}$$

Using Eqs. (11) and (48) and making the corresponding parameter replacements in Eqs. (24) and (28), we obtained energy eigenvalues equation and eigenfunction respectively as

$$E_n=\frac{\alpha^2}{4}\left[\frac{\frac{V_1-2qV_4-qV_5}{q\alpha^2}-\frac{V_2}{q^2\alpha^2}}{2n+1-\eta\sqrt{1+\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}}-\left(2n+1-\eta\sqrt{1+\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)\right]^2-V_4$$

$$n\geq 0,q\geq 1 \tag{49}$$

$$\Psi_n\left(z\right)=D_nz^{\sqrt{\xi-\beta_4}}(1-qz)^{\frac{1}{2}\left(1-\eta\sqrt{1+\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)}P_n^{\left(2\sqrt{\xi-\beta_4},-\eta\sqrt{1+\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)}(1-2qz) \tag{50}$$

Using the above Eq. (50), we can consequently obtain the normalization constant D_n as

$$D_n=\sqrt{\frac{B_{nq}(p,r)}{A_{nq}\left(p,r\right)}} \tag{51}$$

Where

$$A_{nq}(p,r)=\frac{(-1)^n[\Gamma\left(n+2c+1\right)]^2\Gamma(n+2d+1)}{\Gamma\left(n+2c-p+1\right)\Gamma\left(r+2d+1\right)\Gamma(2c+2d+1)}\sum_{n}^{P=0}\frac{(-1)^{p+r}q^{n-p+r}\Gamma(n+2c+2d+r+1)}{p!r!\left(n-p\right)!\left(n-r\right)!\Gamma(p+2d+1)} \tag{52}$$

$$\left(B_{nq}\left(p,r\right)\right)^{-1}=\frac{1}{n+r-p+2\delta+1}{}_2F_1\left(n+r-p+2\delta+1,\eta H-1-p;n+r-p+2\delta+1;q\right) \tag{53}$$

$$c=\delta,=\sqrt{\xi-\beta_4},\beta_4=\frac{V_4}{4\alpha^2},d=-\frac{\eta H}{2}H=\sqrt{1+\frac{V_2}{q^2\alpha^2}-4\frac{(V_3+V_6)}{q\alpha^2}-3\frac{V_5}{\alpha^2}}$$

6 Discussion

In this section, in the framework of the Schrödinger equation the energy eigenvalues and the corresponding wave functions for the Woods-Saxon potential, Rosen-Morse potential and symmetrical double well potential are obtained by setting appropriate parameters in Eq (12) to zero. The PT-symmetric and the non PT-symmetric solution of this potential are also considered

6.1 Woods-Saxon Potential

On putting $V_3 = V_4 = V_5 = V_6 = 0$, the potential (12) turns to the Woods-Saxon [1, 2, 3, 4]

$$V\left(x\right) = \frac{-V_1 e^{-2\alpha x}}{1+qe^{-2\alpha x}} + \frac{-V_2 e^{-4\alpha x}}{1+qe^{-2\alpha x}} \tag{54}$$

The energy eigenvalues equation and wave function are respectively obtained as

$$E_n = -\frac{\alpha^2}{4}\left[\frac{\frac{V_1}{q\alpha^2}+\frac{V_2}{q^2\alpha^2}}{2n+1-\sqrt{\frac{V_2}{q^2\alpha^2}+1}} - \left(2n+1-\sqrt{\frac{V_2}{q^2\alpha^2}+1}\right)\right]^2, \qquad n \geq 0, q \geq 1, \quad \eta = 1 \tag{55}$$

$$\Psi_n\left(z\right) = P_n z^{\sqrt{\xi}}(1-qz)^{\frac{1}{2}\left(1-\sqrt{\frac{V_2}{q^2\alpha^2}+1}\right)} P_n^{\left(2\sqrt{\xi},-\sqrt{\frac{V_2}{q^2\alpha^2}+1}\right)}(1-2qz) \tag{56}$$

This is consistent with [16]

Using the above Eq.(56), the normalization constant P_n is obtained as

$$P_n = \sqrt{\frac{B_{nq}(p,r)}{A_{nq}\left(p,r\right)}} \tag{57}$$

Where

$$A_{nq}(p,r) = \frac{(-1)^n[\Gamma\left(n+2c+1\right)]^2\Gamma(n+2d+1)}{\Gamma\left(n+2c-p+1\right)\Gamma\left(r+2d+1\right)\Gamma(2c+2d+1)}\sum_{n}^{p=0}\frac{(-1)^{p+r}q^{n-p+r}\Gamma(n+2c+2d+r+1)}{p!r!\left(n-p\right)!\left(n-r\right)!\Gamma(p+2d+1)} \tag{58}$$

$$\left(B_{nq}\left(p,r\right)\right)^{-1} = \frac{1}{n+r-p+2\delta+1}{}_2F_1\left(n+r-p+2\delta+1, \eta H-1-p; n+r-p+2\delta+1; q\right) \tag{59}$$

$$c = \delta = \sqrt{\xi}, \quad d = -\frac{\eta H}{2}, \quad H = \sqrt{\frac{V_2}{q^2\alpha^2+1}}$$

Now, let consider the case in which α is a pure imaginary parameter, i.e. $\alpha \to i\alpha$. Under this replacement and after some manipulations, the potential takes the form

$$V\left(x\right) = \frac{-V_1\left[1+q\left(\cos 2\alpha x - isin2\alpha x\right)\right] + V_2\left[\cos 2\alpha x - isin2\alpha x\right]}{\left(1+q^2\right)\cos 2\alpha x + \left(1-q^2\right)isin2\alpha x + 2q} \tag{60}$$

We obtain the real positive energy eigenvalues as

$$E_n = \frac{\alpha^2}{4}\left[\frac{\frac{V_1}{q\alpha^2}-\frac{V_2}{q^2\alpha^2}}{2n+1-\sqrt{1-\frac{V_2}{q^2\alpha^2}}} - \left(2n+1-\sqrt{1-\frac{V_2}{q^2\alpha^2}}\right)\right]^2, \ \eta = 1, \quad n \geq 0, \quad q \geq 1 \tag{61}$$

The above Eq. (61) is consistent with [16] and [17] for $\hbar = 2\mu = 1$ and $\alpha = \frac{\alpha 1}{2}$. By assuming some of the parameters as pure imaginary, i.e. $q \to iq$, $\alpha \to i\alpha$ and $V_1 \to iV_1$, then we have potential of the form

$$V(x) = \frac{V_1 \left[qcos\alpha x + i\left(1 + qcos2\alpha x\right)\right] + V_2[\cos 2\alpha x - isin2\alpha x]}{\left(1 - q^2\right)\cos 2\alpha x + i\left(1 + q^2\right)\sin 2\alpha x + 2iq} \tag{62}$$

We obtained the corresponding energy eigenvalues as

$$E_n = \frac{\alpha^2}{4}\left[\frac{\frac{V_1}{q\alpha^2} - \frac{V_2}{q^2\alpha^2}}{2n + 1 - \sqrt{\frac{V_2}{q^2\alpha^2} + 1}} - \left(2n + 1 - \sqrt{\frac{V_2}{q^2\alpha^2} + 1}\right)\right]^2, \quad n \geq 0, \quad q \geq 1 \tag{63}$$

6.2 Rosen-Morse Potential

If we substitute for $V_1 = V_2 = V_5 = V_6 = 0$, then the potential in Eq. (12) becomes Rosen-Morse

$$V(x) = -V_3 sech_q^2(\alpha x) - V_4 \tanh_q(\alpha x) \tag{64}$$

The energy eigenvalues equation and corresponding wave function are respectively obtained as

$$E_n = -\frac{\alpha^2}{4}\left[\frac{\frac{2qV_4}{q\alpha^2}}{2n + 1 + \sqrt{1 + 4\frac{V_3}{q\alpha^2}}} - \left(2n + 1 + \sqrt{1 + 4\frac{V_3}{q\alpha^2}}\right)\right]^2 - V_4$$

$$= -\frac{\alpha^2}{4}\left(2n + 1 + \sqrt{1 + 4\frac{V_3}{q\alpha^2}}\right)^2 - \frac{V_4^2}{\alpha^2}\left(2n + 1 + \sqrt{1 + 4\frac{V_3}{q\alpha^2}}\right)^{-2}, \quad n \geq 0, \quad q \geq 1 \tag{65}$$

$$\Psi_n(z) = R_n z^{\sqrt{\xi - \beta_4}}(1 - qz)^{\frac{1}{2}\left(1 + \sqrt{1 + 4\frac{V_3}{q\alpha^2}}\right)} P_n^{\left(2\sqrt{\xi - \beta_4}, \sqrt{1 + 4\frac{V_3}{q\alpha^2}}\right)}(1 - 2qz), \ \eta = -1 \tag{66}$$

Which is consistent with [10] Using the above Eq. (66), we obtained the normalization constant R_n as

$$R_n = \sqrt{\frac{B_{nq}(p, r)}{A_{nq}(p, r)}} \tag{67}$$

Where

$$A_{nq}(p, r) = \frac{(-1)^n \left[\Gamma\left(n + 2c + 1\right)\right]^2 \Gamma(n + 2d + 1)}{\Gamma\left(n + 2c - p + 1\right)\Gamma\left(r + 2d + 1\right)\Gamma(2c + 2d + 1)} \sum_{n}^{P=0} \frac{(-1)^{p+r} q^{n-p+r}\Gamma(n + 2c + 2d + r + 1)}{p!r!\left(n - p\right)!\left(n - r\right)!\Gamma(p + 2d + 1)} \tag{68}$$

$$\left(B_{nq}\left(p, r\right)\right)^{-1} = \frac{1}{n + r - p + 2\delta + 1}{}_2F_1\left(n + r - p + 2\delta + 1, \eta H - 1 - p; n + r - p + 2\delta + 1; q\right) \tag{69}$$

$$c = \delta = \sqrt{\xi - \beta_4}, \quad d = -\frac{\eta H}{2}, \quad H = \sqrt{\frac{V_3}{q\alpha^2} + 1}$$

We consider a case in which α is a pure imaginary parameter, i.e. $\alpha \to i\alpha$. Then Eq. (64) takes the form

$$V\left(x\right) = \frac{-4V_3 - V_4\left[\left(1-q^2\right)\cos 2\alpha x + i\left(1+q^2\right)\sin 2\alpha x\right]}{\left(1+q^2\right)\cos 2\alpha x + \left(1-q^2\right)isin2\alpha x + 2q} \tag{70}$$

The positive energy eigenvalues is given as

$$E_n = \frac{\alpha^2}{4}\left[\frac{\frac{2V_4}{\alpha^2}}{2n+1-\sqrt{1-4\frac{V_3}{q\alpha^2}}} - \left(2n+1-\sqrt{1-4\frac{V_3}{q\alpha^2}}\right)\right]^2 - V_4, \quad n \geq 0, q \geq 1 \tag{71}$$

Setting some of the parameters as pure imaginary, i.e. $q \to iq$,$\alpha \to i\alpha$, $V_3 \to iV_3$ and $V_4 \to iV_4$, Then Eq. (64) takes the form

$$V\left(x\right) = \frac{-iV_3 - V_4\left[\left(1+q^2\right)\cos 2\alpha x + i(1-q^2)\sin 2\alpha x\right]}{\left(1-q^2\right)\cos 2\alpha x + i\left(1+q^2\right)\sin 2\alpha x + 2iq} \tag{72}$$

The energy eigenvalues equation is also found to be (71)

6.3 Symmetrical Double Well Potential

Choosing $V_1 = V_2 = V_3 = V_4 = 0$ in Eq. (12) we have

$$V\left(x\right) = V_5 \tanh_q^2\left(\alpha x\right) - V_6 sech_q^2(\alpha x) \tag{73}$$

Again, energy eigenvalues and the corresponding wave function are respectively obtained as

$$E_n = -\frac{\alpha^2}{4}\left[\frac{\frac{V_5}{\alpha^2}}{2n+1-\eta\sqrt{4\frac{V_6}{q\alpha^2}+\frac{3V_5}{\alpha^2}+1}} - \left(2n+1-\eta\sqrt{4\frac{V_6}{q\alpha^2}+\frac{3V_5}{\alpha^2}+1}\right)\right]^2, \ n \geq 0, \ q \geq 1 \tag{74}$$

$$\Psi_n\left(z\right) = W_n z^{\sqrt{\xi}}(1-qz)^{\frac{1}{2}\left(1-\eta\sqrt{4\frac{V_6}{q\alpha^2}+\frac{3V_5}{\alpha^2}+1}\right)} P_n^{\left(2\sqrt{\xi-\beta_4}, -\eta\sqrt{4\frac{V_6}{q\alpha^2}+\frac{3V_5}{\alpha^2}+1}\right)}(1-2qz), \eta = \pm 1 \tag{75}$$

Using the above Eq. (75), we obtained the normalization constant W_n as

$$W_n = \sqrt{\frac{B_{nq}(p,r)}{A_{nq}\left(p,r\right)}} \tag{76}$$

Where

$$A_{nq}(p,r) = \frac{(-1)^n\left[\Gamma\left(n+2c+1\right)\right]^2\Gamma(n+2d+1)}{\Gamma\left(n+2c-p+1\right)\Gamma\left(r+2d+1\right)\Gamma(2c+2d+1)}\sum_{n}^{P=0}\frac{(-1)^{p+r}q^{n-p+r}\Gamma(n+2c+2d+r+1)}{p!r!\left(n-p\right)!\left(n-r\right)!\Gamma(p+2d+1)} \tag{77}$$

$$(B_{nq}(p,r))^{-1} = \frac{1}{n+r-p+2\delta+1}\,{}_2F_1\left(n+r-p+2\delta+1, \eta H-1-p; n+r-p+2\delta+1; q\right) \tag{78}$$

$$c = \delta = \sqrt{\xi}, \quad d = -\frac{\eta H}{2}, \quad H = \sqrt{4\frac{V_6}{q\alpha^2} + \frac{3V_5}{\alpha^2} + 1}$$

Let us consider the case where at least one of the potential parameters is complex. If α is a pure imaginary parameter, i.e. $\alpha \to i\alpha$, symmetrical double well potential can be written as

$$V(x) = \frac{-4V_6 + V_5\left[(1+q^2)\cos 2\alpha x + i(1-q^2)\sin 2\alpha x - 2q\right]}{(1+q^2)\cos 2\alpha x + (1-q^2)\, isin 2\alpha x + 2q} \tag{79}$$

By using Eqs. (11) and (78) and making the corresponding parameter replacements in Eq. (24) again, we obtained the real positive energy eigenvalues equation

$$E_n = \frac{\alpha^2}{4}\left[\frac{\frac{qV_5}{q\alpha^2}}{2n+1-\eta\sqrt{1-4\frac{V_6}{q\alpha^2}-3\frac{V_5}{\alpha^2}}} - \left(2n+1-\eta\sqrt{1-4\frac{V_6}{q\alpha^2}-3\frac{V_5}{\alpha^2}}\right)\right]^2, \quad n \geq 0, q \geq 1 \tag{80}$$

When $\alpha \to i\alpha$, $V_6 \to iV_6$ and $q, \in \Im$ then $V(x)$ takes the form

$$V(x) = \frac{-iV_6 + V_5\left[(1+q^2)\cos 2\alpha x + i(1-q^2)\sin 2\alpha x - 2iq\right]}{(1-q^2)\cos 2\alpha x + i(1+q^2)\sin 2\alpha x + 2iq} \tag{81}$$

The positive energy eigenvalues is also found to be (80)

7 Conclusion

In this paper we solve the Schrödinger equation for Woods-Saxon plus Rosen-Morse plus Symmetrical double well potential via Nikiforov-Uvarov mathematical method. The results include energy spectrum of Wood-Saxon potential, Rosen-Morse potential and Symmetrical double well potential which are obtained by setting some parameters to zero. The PT-symmetric and non-PT-symmetric of Wood-Saxon potential, Rosen-Morse potential symmetrical double well potential have also been discussed.

References

[1] H. Fakhri, and J. Sadeghi, Mod. Phys. Lett. A **19**, 615 (2004).

[2] C. Berkdemir, A. Berkdemir and R. Sever, J. Math. Chem. **43**, 944 (2008).

[3] C. Berkdemir, A. Berkdemir and R. Sever, J. Phys. A: Math. Gen. **39**, 13455 (2006).

[4] C. Berkdemir, A. Berkdemir and R. Sever, Phys. Rev. C **72** 027001 (2005).

[5] Z. Ahmed, Phys. Lett. A **294** 287 (2002).

[6] X. Y. Gu and J. Q. Sun, J.Math. Phys. 51, 022106-022111 (2010).

[7] M. Simsek and H. Erifes, J. Phys. A: Math. Gen. **37**, 4379 (2004).

[8] S. Meyur and S. Debnath, Mod. Phys. Lett. A **23**, 2077 (2008).

[9] M. Q. Zhong, A. Gonzalez-Cisneros, B. W. Xu and S. H. Dong, Phy. Lett. A. **371** 180 (2007)

[10] A. R. Akbarieh and H. Motavalli, Mod. Phys. Lett. A. **23** 3005 (2008).

[11] S. Meyur and S. Debnath, Lat. Am. J. Phys. Educ. **3** 300 (2009).

[12] A. F. Nikiforov and V. B. Uvarov, *Special Functions of Mathematical Physics*, (Basel, Birkhauser) (1988).

[13] A. F. Nikiforov, S. K. Suslor and V. B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*, Springer, Berlin, (1991).

[14] A. Arai, J. Math. Anal. Appl. **158** 63 (1991).

[15] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical functions with formulas, Graphs and Mathematical Tables* (Dover, New York) (1970).

[16] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, (Academic Press, New York) (2007).

[17] S. Meyur and S. Debnath, Lat. Am. J. Phys. Educ. **4** 3 2010.

[18] S. M. Ikhdair and R. Sever, Int. J. Theo. Phys. **46** 1643 (2007).